



Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Games and Economic Behavior 49 (2004) 260–287

**GAMES and
Economic
Behavior**

www.elsevier.com/locate/geb

Best response equivalence

Stephen Morris^{a,*}, Takashi Ui^b

^a *Department of Economics, Yale University, USA*

^b *Faculty of Economics, Yokohama National University, Japan*

Received 2 August 2002

Available online 15 April 2004

Abstract

Two games are best-response equivalent if they have the same best-response correspondence. We provide a characterization of when two games are best-response equivalent. The characterizations exploit a dual relationship between payoff differences and beliefs. Some “potential game” arguments [*Games Econ. Behav.* 14 (1996) 124] rely only on the property that potential games are best-response equivalent to identical interest games. Our results show that a large class of games are best-response equivalent to identical interest games, but are not potential games. Thus we show how some existing potential game arguments can be extended.

© 2004 Elsevier Inc. All rights reserved.

JEL classification: C72

Keywords: Best response equivalence; Duality; Farkas’ Lemma; Potential games

1. Introduction

We consider three progressively stronger equivalence relations on games and characterize each of them.

- Two games are best-response equivalent if they have the same best-response correspondence.
- Two games are better-response equivalent if, for every pair of strategies, they agree when one strategy is better than the other.

* Corresponding author’s address: Cowles Foundation, P.O. Box 208281, New Haven, CT 06520-8281, USA.
E-mail addresses: stephen.morris@yale.edu (S. Morris), oui@ynu.ac.jp (T. Ui).

- Two games are von Neumann–Morgenstern equivalent (VNM-equivalent) if, for each player, the payoff function in one game is equal to a constant times the payoff function in the other game, plus a function that depends only on the opponents' strategies.

Two games are VNM-equivalent if and only if, for each player i , there is a constant $w_i > 0$ such that the ratio of payoff differences from switching between one strategy to another strategy is always w_i . The constant w_i is thus independent of the strategies being compared.

Two games are better-response equivalent if and only if they have the same dominance relations and, for each player i and each pair of strategies a_i and a'_i such that neither strategy strictly dominates the other, there exists a constant $w_i > 0$ such that the ratio of payoff differences from switching between a_i and a'_i is always w_i . In general, this is a weaker requirement than VNM-equivalence. It is weaker both because the proportional payoff differences property is no longer required to hold between some strategy pairs, and because the weight w_i is not necessarily independent of the strategy pair. But if the game does not have dominated strategies, the weights can no longer depend on the strategies being compared, and better-response equivalence collapses to VNM-equivalence.

Two games are best-response equivalent if and only if, for each player i and each pair of strategies a_i and a'_i such that both strategies are a best response to some belief, there exists a constant $w_i > 0$ such that the ratio of payoff differences from switching between a_i and a'_i is always w_i . Even if a game has no dominated strategies, this is a weaker requirement than VNM-equivalence. In games with diminishing marginal returns, best-response equivalence is always a strictly weaker requirement than VNM-equivalence. Examples are given in the paper.

The most extensive discussion and applications of these relations have come in the literature on potential games. Monderer and Shapley (1996b) said that a game was a “potential game” if there exists a potential function, defined on the strategy space, with the property that the change in any player's payoff function from switching between any two of his strategies (holding other players' strategies fixed) was equal to the change in the potential function.¹ A game is “weighted potential game,” if the payoff changes are proportional for each player. Thus a game is a weighted potential game if and only if it is VNM-equivalent to a game with identical payoff functions. While some results using potential or weighted potential game arguments are using the VNM-equivalence to identical interest games, other arguments are just using the better-response equivalence and even only best-response equivalence implications of VNM-equivalence.² Any paper that deals only with equilibria is using only best-response equivalence (e.g., Neyman, 1997; Ui, 2001; Morris and Ui, 2002). Similarly, fictitious play only uses the best-response

¹ See also Ui (2000) for a characterization and examples of potential games.

² Arguments that exploit potential arguments to prove the existence of a pure strategy equilibrium (e.g., Rosenthal, 1973) only use *ordinal* properties of payoffs. Monderer and Shapley (1996b) introduced ordinal potential games and Voorneveld (2000) and Dubey et al. (2002) showed how ordinal potential games can be weakened to only require pure strategy best-response equivalence.

properties of the game (Monderer and Shapley, 1996a).³ An application using only better-response equivalence but not the VNM-equivalence appears in Morris (1999). Some papers studying quantal responses or stochastic best responses in potential games use the full power of VNM-equivalence (e.g., Blume, 1993; Brock and Durlauf, 2001; Anderson et al., 2001; Ui, 2002).⁴

The fact that VNM-equivalence is the same as better-response equivalence in the absence of dominated strategies and may be different in the presence of dominated strategies has been noted in a number of contexts (see Sela, 1992; Blume, 1993, p. 409; Monderer and Shapley, 1996b, footnote 9; Maskin and Tirole, 2001, p. 209). However, our characterizations of better-response equivalence in the presence of dominated strategies and of the significant gap between better-response equivalence and best-response equivalence fill a gap in the literature.⁵

The paper is organized as follows. In Section 2, we describe our notions of equivalence and give an example illustrating the differences. In Section 3, we report our characterizations. In Section 4, we restrict attention to a class of games where best-response equivalence is a strictly weaker requirement than VNM-equivalence and characterize that class. We also discuss an extension to games with infinite strategy spaces and its application. Section 5 briefly discusses better-response and best-response equivalence in the mixed strategy extension of a game.

2. Equivalence properties of games

A game consists of a finite set of players N and a finite strategy set A_i for $i \in N$, and a payoff function $g_i : A \rightarrow \mathbb{R}$ for $i \in N$ where $A = \prod_{i \in N} A_i$. We write $A_{-i} = \prod_{j \neq i} A_j$ and $a_{-i} = (a_j)_{j \neq i} \in A_{-i}$. We simply denote a game by $\mathbf{g} = (g_i)_{i \in N}$. Throughout the paper, we regard $g_i(a_i, \cdot) : A_{-i} \rightarrow \mathbb{R}$ as a vector in $\mathbb{R}^{A_{-i}}$. We write $g_i(a_i, \cdot) \gg g_i(a'_i, \cdot)$ if $g_i(a_i, a_{-i}) > g_i(a'_i, a_{-i})$ for all $a_{-i} \in A_{-i}$, and $g_i(a_i, \cdot) \geq g_i(a'_i, \cdot)$ if $g_i(a_i, a_{-i}) \geq g_i(a'_i, a_{-i})$ for all $a_{-i} \in A_{-i}$.

For $i \in N$, let $\Delta(A_{-i})$ denote the set of all probability distributions over A_{-i} . We call each element of $\Delta(A_{-i})$ player i 's belief. For $X_i \subseteq A_i$, let $\Lambda_i(a_i, X_i | g_i) \subseteq \Delta(A_{-i})$ be a set of player i 's beliefs such that player i with a payoff function g_i and a belief $\lambda_i \in \Lambda_i(a_i, X_i | g_i)$ weakly prefers a_i to any strategy in X_i :

$$\begin{aligned} & \Lambda_i(a_i, X_i | g_i) \\ &= \left\{ \lambda_i \in \Delta(A_{-i}) \mid \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i}) (g_i(a_i, a_{-i}) - g_i(a'_i, a_{-i})) \geq 0 \text{ for all } a'_i \in X_i \right\}. \end{aligned}$$

When X_i is a singleton, i.e., $X_i = \{a'_i\}$, we write $\Lambda_i(a_i, a'_i | g_i)$ instead of $\Lambda_i(a_i, \{a'_i\} | g_i)$.

³ Sela (1999) establishes convergence of fictitious play in a class of "one-against-all" games. These games are best-response equivalent to identical interest games, but not potential games.

⁴ More precisely, they use the full power of VNM-equivalence such that the constant w_i is the same for all the players.

⁵ Mertens (1987) studied various notions of best-response equivalence, but with his more abstract strategy spaces and focus on admissible best responses, there is little overlap with the material in this paper.

We are interested in characterizing two equivalence relations on games captured by these sets of beliefs by which players prefer one particular strategy.

Definition 1. A game \mathbf{g} is *better-response equivalent* to $\mathbf{g}' = (g'_i)_{i \in N}$ if, for each $i \in N$,

$$\Lambda_i(a_i, a'_i | g_i) = \Lambda_i(a_i, a'_i | g'_i)$$

for all $a_i, a'_i \in A_i$.

Definition 2. A game \mathbf{g} is *best-response equivalent* to $\mathbf{g}' = (g'_i)_{i \in N}$ if, for each $i \in N$,

$$\Lambda_i(a_i, A_i | g_i) = \Lambda_i(a_i, A_i | g'_i)$$

for all $a_i \in A_i$.

If \mathbf{g} is better-response equivalent to \mathbf{g}' , then \mathbf{g} is best-response equivalent to \mathbf{g}' , since

$$\Lambda_i(a_i, A_i | g_i) = \bigcap_{a'_i \in A_i} \Lambda_i(a_i, a'_i | g_i).$$

An easy sufficient condition for better-response equivalence is the following.⁶

Definition 3. A game \mathbf{g} is *VNM-equivalent* to $\mathbf{g}' = (g'_i)_{i \in N}$ if, for each $i \in N$, there exists a positive constant $w_i > 0$ and a function $Q_i : A_{-i} \rightarrow \mathbb{R}$ such that

$$g_i(a_i, \cdot) = w_i g'_i(a_i, \cdot) + Q_i(\cdot)$$

for all $a_i \in A_i$.

It is straightforward to see that if \mathbf{g} is VNM-equivalent to \mathbf{g}' , then

$$g_i(a_i, \cdot) - g_i(a'_i, \cdot) = w_i (g'_i(a_i, \cdot) - g'_i(a'_i, \cdot))$$

for all $a_i, a'_i \in A_i$. Conversely, if this is true, then a function $Q_i : A_{-i} \rightarrow \mathbb{R}$ such that

$$Q_i(\cdot) = g_i(a_i, \cdot) - w_i g'_i(a_i, \cdot)$$

is well defined, and thus \mathbf{g} is VNM-equivalent to \mathbf{g}' . Thus, we have the following lemma.

Lemma 1. A game \mathbf{g} is VNM-equivalent to \mathbf{g}' if and only if, for each $i \in N$, there exists w_i such that

$$g_i(a_i, \cdot) - g_i(a'_i, \cdot) = w_i (g'_i(a_i, \cdot) - g'_i(a'_i, \cdot)) \quad (1)$$

for all $a_i, a'_i \in A_i$.

It is straightforward to see that VNM-equivalence is sufficient for better-response equivalence. In fact, (1) implies that

⁶ Blume (1993) called this property “strongly best-response equivalent.”

$$\begin{aligned} & \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(a_i, a_{-i}) - g_i(a'_i, a_{-i})) \\ &= w_i \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g'_i(a_i, a_{-i}) - g'_i(a'_i, a_{-i})) \end{aligned}$$

for all $\lambda_i \in \Delta(A_{-i})$ and thus $\Lambda_i(a_i, a'_i | g_i) = \Lambda_i(a_i, a'_i | g'_i)$ for all $a_i, a'_i \in A_i$.

Best-response, better-response, and VNM-equivalence are equivalence relations. Thus, they define an equivalence class of games. For example, weighted potential games (Monderer and Shapley, 1996b) with a weighted potential function $f : A \rightarrow \mathbb{R}$ are regarded as a VNM-equivalence class of an identical interest game $\mathbf{f} = (f_i)_{i \in N}$ with $f_i = f$ for all $i \in N$. This is clear by Lemma 1 and the following original definition of weighted potential games.

Definition 4. A game $\mathbf{g} = (g_i)_{i \in N}$ is a *weighted potential game* if there exists a weighted potential function $f : A \rightarrow \mathbb{R}$ and $w_i > 0$ for each $i \in N$ such that

$$g_i(a_i, \cdot) - g_i(a'_i, \cdot) = w_i(f(a_i, \cdot) - f(a'_i, \cdot))$$

for all $a_i, a'_i \in A_i$. If $w_i = 1$ for all $i \in N$, \mathbf{g} is called a potential game and f is called a potential function.

As the concept of VNM-equivalence leads us to the definition of weighted potential games, the concept of better-response equivalence and that of best-response equivalence lead us to the following definitions of new classes of games.

Definition 5. A game $\mathbf{g} = (g_i)_{i \in N}$ is a *better-response potential game* if it is better-response equivalent to an identical interest game $\mathbf{f} = (f_i)_{i \in N}$ with $f_i = f$ for all $i \in N$. A function f is called a better-response potential function.

Definition 6. A game $\mathbf{g} = (g_i)_{i \in N}$ is a *best-response potential game* if it is best-response equivalent to an identical interest game $\mathbf{f} = (f_i)_{i \in N}$ with $f_i = f$ for all $i \in N$. A function f is called a best-response potential function.

Voorneveld (2000) called a game a best-response potential game if its best-response correspondence coincides with that of an identical interest game over the class of beliefs such that $\lambda_i(a_{-i}) = 0$ or 1. Thus, best-response potential games in this paper form a special class of those in Voorneveld (2000).

Existing potential game results that rely only on better-response equivalence or best-response equivalence, such as those mentioned in the introduction, automatically hold for the larger class of better-response potential games or that of best-response potential games. Thus, we are interested in exactly when and to what extent better-response and best-response equivalence are weaker requirements than VNM-equivalence.

Notice that best-response and better-response equivalence are clearly weaker requirements than VNM-equivalence, because the latter imposes too many constraints on payoffs from dominated strategies. Moreover, best-response equivalence is significantly weaker than better-response equivalence, as shown by the following example.

Consider a two player, three strategy, symmetric payoff game $\mathbf{g}(x, y)$ parameterized by $(x, y) \in \mathbb{R}_{++}^2$, where each player's payoffs are given by the following payoff matrix (where the player's own strategies are represented by rows and his opponent's strategies are represented by columns):

	1	2	3
1	x	$-x$	$-2x$
2	0	0	0
3	$-2y$	$-y$	y

In the special case where $x = y = 1$, we have game $\mathbf{g}(1, 1)$ with the following payoff matrix:

	1	2	3
1	1	-1	-2
2	0	0	0
3	-2	-1	1

If a row player has a belief $\lambda_i(k) = \pi_k$ for $k \in \{1, 2, 3\}$, he prefers strategy 1 to strategy 2 if and only if

$$\pi_1 \geq \pi_2 + 2\pi_3;$$

he prefers strategy 1 to strategy 3 if and only if

$$(x + 2y)\pi_1 \geq (x - y)\pi_2 + (2x + y)\pi_3;$$

he prefers strategy 3 to strategy 2 if and only if

$$\pi_3 \geq \pi_2 + 2\pi_1.$$

Thus the region of indifference between strategies 1 and 2, and between strategy 2 and 3, does not depend on x and y . Moreover, whenever strategy 1 (or 3) is preferred to strategy 2, it is also preferred to strategy 3 (or 1). Thus the best response regions for this game are as in Fig. 1, for any $(x, y) \in \mathbb{R}_{++}^2$. Thus $\mathbf{g}(x, y)$ is best-response equivalent to $\mathbf{g}(1, 1)$ for any




Fig. 1. The best response regions.

$(x, y) \in \mathbb{R}_{++}^2$. On the other hand, the region of indifference between strategies 1 and 3 does depend on x and y : in particular, $\mathbf{g}(x, y)$ is better-response equivalent to $\mathbf{g}(1, 1)$ if and only if $x = y$. We will discuss this example again in Section 4.

3. Results

3.1. Generic properties of games

We will appeal to some generic properties of games, i.e., properties that will hold for all but a Lebesgue measure zero set of payoffs (as long as each player has at least two actions).

G1. For all $i \in N$, if $g_i(a_i, \cdot) \geq g_i(a'_i, \cdot)$, then $g_i(a_i, \cdot) \gg g_i(a'_i, \cdot)$ for distinct $a_i, a'_i \in A_i$.

G2. For all $i \in N$, vectors $g_i(a_i, \cdot) - g_i(a'_i, \cdot)$ and $g_i(a_i, \cdot) - g_i(a''_i, \cdot)$ are linearly independent for distinct $a_i, a'_i, a''_i \in A_i$.

G3. For all $i \in N$, if $\Lambda_i(a_i, A_i | g_i) \cap \Lambda_i(a'_i, A_i | g_i) \neq \emptyset$, then

$$\Lambda_i(a_i, A_i \setminus \{a'_i\} | g_i) \setminus \Lambda_i(a_i, a'_i | g_i) \neq \emptyset \quad \text{for distinct } a_i, a'_i \in A_i.$$

3.2. Better-response equivalence

Strategy a_i strictly dominates a'_i in game \mathbf{g} (we write $a_i \succ_i^{\mathbf{g}} a'_i$) if $g_i(a_i, \cdot) \gg g_i(a'_i, \cdot)$, or, equivalently, $\Lambda_i(a'_i, a_i | g_i) = \emptyset$. Strategies a_i and a'_i are better-response comparable (we write $a_i \sim_i^{\mathbf{g}} a'_i$) if neither $a_i \succ_i^{\mathbf{g}} a'_i$ nor $a'_i \succ_i^{\mathbf{g}} a_i$.

Proposition 1. If games \mathbf{g} and \mathbf{g}' satisfy generic property G1, then \mathbf{g} is better-response equivalent to \mathbf{g}' if and only if, for each $i \in N$, (a) they have the same dominance relations ($\succ_i^{\mathbf{g}} = \succ_i^{\mathbf{g}'}$) and (b) whenever a_i is better-response comparable to a'_i ($a_i \sim_i^{\mathbf{g}} a'_i$), there exists $w_i(a_i, a'_i) > 0$ such that

$$g_i(a_i, \cdot) - g_i(a'_i, \cdot) = w_i(a_i, a'_i)(g'_i(a_i, \cdot) - g'_i(a'_i, \cdot)). \quad (2)$$

Farkas' Lemma⁷ plays a central role in the proofs.

Lemma 2 (Farkas' Lemma). For vectors $\mathbf{a}_0, \mathbf{a}_1, \dots, \mathbf{a}_m \in \mathbb{R}^n$, the following two conditions are equivalent.

- If $(\mathbf{a}_1 \cdot \mathbf{y}), \dots, (\mathbf{a}_m \cdot \mathbf{y}) \leq 0$ for $\mathbf{y} \in \mathbb{R}^n$, then $(\mathbf{a}_0 \cdot \mathbf{y}) \leq 0$.⁸
- There exists $x_1, \dots, x_m \geq 0$ such that $x_1 \mathbf{a}_1 + \dots + x_m \mathbf{a}_m = \mathbf{a}_0$.

⁷ See a textbook of convex analysis such as the recent one by Hiriart-Urruty and Lemaréchal (2001), or the classic one by Rockafellar (1970).

⁸ I.e., if $\sum_{j=1}^n a_{ij} y_j \leq 0$ for each $i = 1, \dots, m$, then $\sum_{j=1}^n a_{0j} y_j \leq 0$.

Proof of Proposition 1. We first show that (a) and (b) are sufficient for the better-response equivalence of \mathbf{g} and \mathbf{g}' . If $a_i \sim_i^{\mathbf{g}} a'_i$, then (b) implies that

$$\begin{aligned} & \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(a_i, a_{-i}) - g_i(a'_i, a_{-i})) \\ &= w_i(a_i, a'_i) \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g'_i(a_i, a_{-i}) - g'_i(a'_i, a_{-i})) \end{aligned}$$

and thus

$$\Lambda_i(a_i, a'_i | g_i) = \Lambda_i(a_i, a'_i | g'_i).$$

If $a_i \succ_i^{\mathbf{g}} a'_i$, then

$$\Lambda_i(a_i, a'_i | g_i) = \Lambda_i(a_i, a'_i | g'_i) = \Delta(A_{-i}).$$

If $a'_i \succ_i^{\mathbf{g}} a_i$, then

$$\Lambda_i(a_i, a'_i | g_i) = \Lambda_i(a_i, a'_i | g'_i) = \emptyset.$$

To prove necessity, suppose that \mathbf{g} is better-response equivalent to \mathbf{g}' . Since

$$\Lambda_i(a_i, a'_i | g_i) = \Lambda_i(a_i, a'_i | g'_i),$$

we have

$$a_i \succ_i^{\mathbf{g}} a'_i \Leftrightarrow \Lambda_i(a'_i, a_i | g_i) = \Lambda_i(a'_i, a_i | g'_i) = \emptyset \Leftrightarrow a_i \succ_i^{\mathbf{g}'} a'_i$$

and thus (a) holds.

To prove (b), suppose that $a_i \sim_i^{\mathbf{g}} a'_i$. We know that $a_i \sim_i^{\mathbf{g}'} a'_i$. Let $\lambda_i \in \Delta(A_{-i})$ be such that

$$\sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(a_i, a_{-i}) - g_i(a'_i, a_{-i})) \geq 0.$$

Since $\lambda_i \in \Lambda_i(a_i, a'_i | g_i) = \Lambda_i(a_i, a'_i | g'_i)$,

$$\sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g'_i(a_i, a_{-i}) - g'_i(a'_i, a_{-i})) \geq 0.$$

This implies that if $(y_{a_{-i}})_{a_{-i} \in A_{-i}} \in \mathbb{R}^{A_{-i}}$ is such that

$$\begin{aligned} & - \sum_{a_{-i} \in A_{-i}} y_{a_{-i}}(g_i(a_i, a_{-i}) - g_i(a'_i, a_{-i})) \leq 0, \\ & -y_{a_{-i}} \leq 0 \quad \text{for all } a_{-i} \in A_{-i}, \end{aligned}$$

then

$$- \sum_{a_{-i} \in A_{-i}} y_{a_{-i}}(g'_i(a_i, a_{-i}) - g'_i(a'_i, a_{-i})) \leq 0.$$

By Farkas' Lemma, there exist $x_{a'_i}^{a_i} \geq 0$ and $z_{a_{-i}} \geq 0$ for $a_{-i} \in A_{-i}$ such that

$$-x_{a'_i}^{a_i}(g_i(a_i, \cdot) - g_i(a'_i, \cdot)) - \sum_{a_{-i} \in A_{-i}} z_{a_{-i}} \delta^{a_{-i}}(\cdot) = -(g'_i(a_i, \cdot) - g'_i(a'_i, \cdot))$$

where $\delta^{a-i} : A_{-i} \rightarrow \mathbb{R}$ is such that $\delta^{a-i}(a'_{-i}) = 1$ if $a'_{-i} = a_{-i}$ and $\delta^{a-i}(a'_{-i}) = 0$ otherwise. Thus,

$$x_{a'_i}^{a_i} (g_i(a_i, \cdot) - g_i(a'_i, \cdot)) \leq g'_i(a_i, \cdot) - g'_i(a'_i, \cdot).$$

If $x_{a'_i}^{a_i} = 0$, then $g'_i(a_i, \cdot) - g'_i(a'_i, \cdot) \geq 0$. However, this is impossible since $a_i \sim_i^{\mathbf{g}'} a'_i$ implies that a_i does not strictly dominate a'_i in \mathbf{g}' and G1 requires that if a_i does not strictly dominate a'_i , then it is not the case that $g'_i(a_i, \cdot) - g'_i(a'_i, \cdot) \geq 0$. Thus, $x_{a'_i}^{a_i} > 0$.

Symmetrically, we have

$$x_{a_i}^{a'_i} (g_i(a'_i, \cdot) - g_i(a_i, \cdot)) \leq g'_i(a'_i, \cdot) - g'_i(a_i, \cdot)$$

where $x_{a_i}^{a'_i} > 0$. Thus,

$$(x_{a'_i}^{a_i} - x_{a_i}^{a'_i}) (g_i(a_i, \cdot) - g_i(a'_i, \cdot)) \leq 0.$$

If $x_{a'_i}^{a_i} - x_{a_i}^{a'_i} > 0$, then $g_i(a_i, \cdot) - g_i(a'_i, \cdot) \leq 0$, and if $x_{a'_i}^{a_i} - x_{a_i}^{a'_i} < 0$, then $g_i(a_i, \cdot) - g_i(a'_i, \cdot) \geq 0$, which we already noted are impossible. Thus, $x_{a'_i}^{a_i} = x_{a_i}^{a'_i}$, which implies that

$$x_{a'_i}^{a_i} (g_i(a_i, \cdot) - g_i(a'_i, \cdot)) = g'_i(a_i, \cdot) - g'_i(a'_i, \cdot).$$

This proves (b). \square

If \mathbf{g} has no dominated strategy, then (2) is true for every $a_i, a'_i \in A_i$. If $w_i(a_i, a'_i)$ is the same for every $a_i, a'_i \in A_i$, then better-response equivalence implies VNM-equivalence. However, Proposition 1 does not say anything about whether $w_i(a_i, a'_i)$ does depend upon $a_i, a'_i \in A_i$. Thus, we are interested in when better-response equivalence implies VNM-equivalence. The following proposition provides a sufficient condition for the equivalence of better-response equivalence and VNM-equivalence.

Proposition 2. *Suppose that games \mathbf{g} and \mathbf{g}' satisfy generic properties G1 and G2, and that, for each $i \in N$ and for any $a_i, a'_i \in A_i$, there exists a sequence $\{a_i^k\}_{k=1}^m$ such that $a_i^1 = a_i$, $a_i^m = a'_i$, $a_i^k \sim_i^{\mathbf{g}} a_i^{k+1}$ for $k = 1, \dots, m-1$, and $a_i^k \sim_i^{\mathbf{g}} a_i^{k+2}$ for $k = 1, \dots, m-2$. Then \mathbf{g} is better-response equivalent to \mathbf{g}' if and only if \mathbf{g} is VNM-equivalent to \mathbf{g}' .*

Note that the above condition concerning $\sim_i^{\mathbf{g}}$ is trivially satisfied if no strategy is dominated, i.e., $\sim_i^{\mathbf{g}}$ is the complete relation. So, the proposition immediately has the following corollary.

Corollary 3. *If \mathbf{g} and \mathbf{g}' satisfy generic properties G1 and G2 and have no strictly dominated strategies, then \mathbf{g} is better-response equivalent to \mathbf{g}' if and only if \mathbf{g} is VNM-equivalent to \mathbf{g}' .*

It should be emphasized that the sufficient condition of Proposition 2 is sometimes satisfied even when there are strictly dominated strategies in the game. For example, consider the following two player game, where only the row player’s payoffs are shown:

	1	2
1	4	1
2	1	3
3	2	2
4	3	0

Consider strategies of the row player. We have $1 \sim_i^g 2$, $2 \sim_i^g 3$, $3 \sim_i^g 4$, $1 \sim_i^g 3$, $2 \sim_i^g 4$ as in Fig. 2, satisfying the condition of Proposition 2, while strategy 1 strictly dominates strategy 4.

To prove the proposition, we use the following lemma.

Lemma 3. Suppose that g and g' satisfy generic property G2. For some $A'_i \subseteq A_i$, if there exists $w_i(a_i, a'_i) > 0$ such that

$$g_i(a_i, \cdot) - g_i(a'_i, \cdot) = w_i(a_i, a'_i)(g'_i(a_i, \cdot) - g'_i(a'_i, \cdot))$$

for all $a_i, a'_i \in A'_i$, then $w_i(a_i, a'_i)$ is the same for all $a_i, a'_i \in A'_i$.

Proof. Without loss of generality, assume that $|A'_i| \geq 3$. For distinct $a_i, b_i, c_i \in A'_i$, there exist $w_i(a_i, b_i), w_i(b_i, c_i), w_i(a_i, c_i) > 0$ such that

$$\begin{aligned} g_i(a_i, \cdot) - g_i(b_i, \cdot) &= w_i(a_i, b_i)(g'_i(a_i, \cdot) - g'_i(b_i, \cdot)), \\ g_i(b_i, \cdot) - g_i(c_i, \cdot) &= w_i(b_i, c_i)(g'_i(b_i, \cdot) - g'_i(c_i, \cdot)), \\ g_i(a_i, \cdot) - g_i(c_i, \cdot) &= w_i(a_i, c_i)(g'_i(a_i, \cdot) - g'_i(c_i, \cdot)). \end{aligned}$$

We first show that $w_i(a_i, b_i) = w_i(b_i, c_i) = w_i(a_i, c_i)$. Since

$$\begin{aligned} &w_i(a_i, b_i)(g'_i(a_i, \cdot) - g'_i(b_i, \cdot)) + w_i(b_i, c_i)(g'_i(b_i, \cdot) - g'_i(c_i, \cdot)) \\ &= g_i(a_i, \cdot) - g_i(b_i, \cdot) + g_i(b_i, \cdot) - g_i(c_i, \cdot) \\ &= g_i(a_i, \cdot) - g_i(c_i, \cdot) \\ &= w_i(a_i, c_i)(g'_i(a_i, \cdot) - g'_i(c_i, \cdot)) \\ &= w_i(a_i, c_i)(g'_i(a_i, \cdot) - g'_i(b_i, \cdot)) + w_i(a_i, c_i)(g'_i(b_i, \cdot) - g'_i(c_i, \cdot)), \end{aligned}$$




Fig. 2. The graph of \sim_i^g .

we have

$$(w_i(a_i, b_i) - w_i(a_i, c_i))(g'_i(a_i, \cdot) - g'_i(b_i, \cdot)) \\ + (w_i(b_i, c_i) - w_i(a_i, c_i))(g'_i(b_i, \cdot) - g'_i(c_i, \cdot)) = 0.$$

By G2, $g'_i(a_i, \cdot) - g'_i(b_i, \cdot)$ and $g'_i(b_i, \cdot) - g'_i(c_i, \cdot)$ are linearly independent and thus it must be true that $w_i(a_i, b_i) = w_i(b_i, c_i) = w_i(a_i, c_i)$.

Similarly, for distinct $b_i, c_i, d_i \in A'_i$, $w_i(b_i, c_i) = w_i(c_i, d_i) = w_i(b_i, d_i)$. Therefore, $w_i(a_i, b_i) = w_i(c_i, d_i)$ for any $a_i, b_i, c_i, d_i \in A'_i$, which completes the proof. \square

We now report the proof of Proposition 2.

Proof of Proposition 2. We show that if \mathbf{g} is better-response equivalent to \mathbf{g}' then \mathbf{g} is VNM-equivalent to \mathbf{g}' . By G1 and Proposition 1, if $a_i \sim_i^{\mathbf{g}} a'_i$, there exist $w_i(a_i, a'_i) > 0$ such that

$$g_i(a_i, \cdot) - g_i(a'_i, \cdot) = w_i(a_i, a'_i)(g'_i(a_i, \cdot) - g'_i(a'_i, \cdot)).$$

If $|A_i| = 2$, this completes the proof by Lemma 1. Suppose that $|A_i| \geq 3$. For $a_i, a'_i \in A_i$, let $\{a_i^k\}_{k=1}^m$ be a sequence such that $a_i^1 = a_i$, $a_i^m = a'_i$, $a_i^k \sim_i^{\mathbf{g}} a_i^{k+1}$ for $k = 1, \dots, m-1$, and $a_i^k \sim_i^{\mathbf{g}} a_i^{k+2}$ for $k = 1, \dots, m-2$. There exist $x_k, y_k > 0$ such that

$$g_i(a_i^k, \cdot) - g_i(a_i^{k+1}, \cdot) = x_k(g'_i(a_i^k, \cdot) - g'_i(a_i^{k+1}, \cdot)), \\ g_i(a_i^{k+1}, \cdot) - g_i(a_i^{k+2}, \cdot) = x_{k+1}(g'_i(a_i^{k+1}, \cdot) - g'_i(a_i^{k+2}, \cdot)), \\ g_i(a_i^k, \cdot) - g_i(a_i^{k+2}, \cdot) = y_k(g'_i(a_i^k, \cdot) - g'_i(a_i^{k+2}, \cdot)).$$

By Lemma 3, $x_k = x_{k+1} = y_k$ for all $k \leq m-2$. By letting $x_k = w_i(a_i, a'_i)$, we have

$$g_i(a_i, \cdot) - g_i(a'_i, \cdot) = \sum_{k=1}^{m-1} (g_i(a_i^k, \cdot) - g_i(a_i^{k+1}, \cdot)) \\ = \sum_{k=1}^{m-1} x_k (g'_i(a_i^k, \cdot) - g'_i(a_i^{k+1}, \cdot)) \\ = w_i(a_i, a'_i)(g'_i(a_i, \cdot) - g'_i(a'_i, \cdot)).$$

To summarize, for all $a_i, a'_i \in A_i$, there exists $w_i(a_i, a'_i) > 0$ satisfying the above equation. By Lemma 3, $w_i(a_i, a'_i)$ is the same for all $a_i, a'_i \in A_i$. By Lemma 1, \mathbf{g} is VNM-equivalent to \mathbf{g}' , which completes the proof. \square

3.3. Best-response equivalence

Strategies a_i and a'_i are *best-response comparable* (we write $a_i \approx_i^{\mathbf{g}} a'_i$) if both strategies are best responses at some belief, i.e., $\Lambda_i(a_i, A_i | g_i) \cap \Lambda_i(a'_i, A_i | g_i) \neq \emptyset$. Note that $a_i \approx_i^{\mathbf{g}} a_i$ if and only if $\Lambda_i(a_i, A_i | g_i) \neq \emptyset$.

Proposition 4. *If games \mathbf{g} and \mathbf{g}' satisfy generic property G3, then \mathbf{g} is best-response equivalent to \mathbf{g}' if and only if, for each $i \in N$, (a) they have the same best-response comparability relation ($\approx_i^{\mathbf{g}} = \approx_i^{\mathbf{g}'}$) and (b) whenever a_i is best-response comparable to a'_i ($a_i \approx_i^{\mathbf{g}} a'_i$), there exists $w_i(a_i, a'_i) > 0$ such that*

$$g_i(a_i, \cdot) - g_i(a'_i, \cdot) = w_i(a_i, a'_i)(g'_i(a_i, \cdot) - g'_i(a'_i, \cdot)).$$

Proof. We first show that (a) and (b) are sufficient for the best-response equivalence of \mathbf{g} and \mathbf{g}' . If $\Lambda_i(a_i, A_i | g_i) = \emptyset$, then $\Lambda_i(a_i, A_i | g_i) = \Lambda_i(a_i, A_i | g'_i) = \emptyset$ because $\Lambda_i(a_i, A_i | g_i) = \emptyset$ implies that $a_i \approx_i^{\mathbf{g}} a_i$ is not true and thus (a) implies that $a_i \approx_i^{\mathbf{g}'} a_i$ is not true. If $\Lambda_i(a_i, A_i | g_i) \neq \emptyset$, then $\{a'_i | a_i \approx_i^{\mathbf{g}} a'_i\} \neq \emptyset$, and we must have

$$\Lambda_i(a_i, A_i | g_i) = \bigcap_{a'_i \in A_i} \Lambda_i(a_i, a'_i | g_i) = \bigcap_{\{a'_i | a_i \approx_i^{\mathbf{g}} a'_i\}} \Lambda_i(a_i, a'_i | g_i). \tag{3}$$

Clearly, (3) is true when $\{a'_i | a_i \approx_i^{\mathbf{g}} a'_i\} = A_i$. To see that (3) is true when $\{a'_i | a_i \approx_i^{\mathbf{g}} a'_i\} \subset A_i$, suppose otherwise. Then,

$$\bigcap_{a'_i \in A_i} \Lambda_i(a_i, a'_i | g_i) \subset \bigcap_{\{a'_i | a_i \approx_i^{\mathbf{g}} a'_i\}} \Lambda_i(a_i, a'_i | g_i),$$

and thus there exists $a''_i \notin \{a'_i | a_i \approx_i^{\mathbf{g}} a'_i\}$ such that

$$\bigcap_{a'_i \in A_i} \Lambda_i(a_i, a'_i | g_i) \subset \bigcap_{a'_i \in A_i \setminus \{a''_i\}} \Lambda_i(a_i, a'_i | g_i).$$

However, this implies that $a_i \approx_i^{\mathbf{g}} a''_i$, which is a contradiction. Thus, (3) must be true. If $a_i \approx_i^{\mathbf{g}} a'_i$, then (b) implies that

$$\begin{aligned} & \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(a_i, a_{-i}) - g_i(a'_i, a_{-i})) \\ &= w_i(a_i, a'_i) \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g'_i(a_i, a_{-i}) - g'_i(a'_i, a_{-i})), \end{aligned}$$

and thus

$$\Lambda_i(a_i, a'_i | g_i) = \Lambda_i(a_i, a'_i | g'_i). \tag{4}$$

Therefore, by (a), (3), and (4), we have $\Lambda_i(a_i, A_i | g_i) = \Lambda_i(a_i, A_i | g'_i)$. This completes the proof of sufficiency.

To prove necessity, suppose that \mathbf{g} is best-response equivalent to \mathbf{g}' . Since

$$\Lambda_i(a_i, A_i | g_i) = \Lambda_i(a_i, A_i | g'_i),$$

we have

$$\Lambda_i(a_i, A_i | g_i) \cap \Lambda_i(a'_i, A_i | g_i) = \Lambda_i(a_i, A_i | g'_i) \cap \Lambda_i(a'_i, A_i | g'_i)$$

and thus $\approx_i^{\mathbf{g}} = \approx_i^{\mathbf{g}'}$. This proves (a).

If $a_i \approx_i^g a'_i$, then there exists $\lambda_i \in \Delta(A_{-i})$ such that

$$\begin{aligned} \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(a_i, a_{-i}) - g_i(a''_i, a_{-i})) &\geq 0 \quad \text{for all } a''_i \in A_i, \\ \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(a'_i, a_{-i}) - g_i(a''_i, a_{-i})) &\geq 0 \quad \text{for all } a''_i \in A_i \setminus \{a_i\}. \end{aligned}$$

Since $\lambda_i \in \Lambda_i(a_i, A_i | g_i) = \Lambda_i(a_i, A_i | g'_i)$,

$$\sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g'_i(a_i, a_{-i}) - g'_i(a'_i, a_{-i})) \geq 0.$$

The above implies that, if $(y_{a_{-i}})_{a_{-i} \in A_{-i}} \in \mathbb{R}^{A_{-i}}$ is such that

$$\begin{aligned} - \sum_{a_{-i} \in A_{-i}} y_{a_{-i}}(g_i(a_i, a_{-i}) - g_i(a'_i, a_{-i})) &\leq 0, \\ - \sum_{a_{-i} \in A_{-i}} y_{a_{-i}}(g_i(a_i, a_{-i}) - g_i(a''_i, a_{-i})) &\leq 0 \quad \text{for all } a''_i \in A_i \setminus \{a_i, a'_i\}, \\ - \sum_{a_{-i} \in A_{-i}} y_{a_{-i}}(g_i(a'_i, a_{-i}) - g_i(a''_i, a_{-i})) &\leq 0 \quad \text{for all } a''_i \in A_i \setminus \{a_i, a'_i\}, \\ -y_{a_{-i}} &\leq 0 \quad \text{for all } a_{-i} \in A_{-i}, \end{aligned}$$

then

$$- \sum_{a_{-i} \in A_{-i}} y_{a_{-i}}(g'_i(a_i, a_{-i}) - g'_i(a'_i, a_{-i})) \leq 0.$$

By Farkas' Lemma, there exist $x_{a'_i}^{a_i} \geq 0$, $\gamma_{a'_i}^{a_i} : A_{-i} \rightarrow \mathbb{R}$, and $\delta_{a'_i}^{a_i} : A_{-i} \rightarrow \mathbb{R}$ such that

$$-x_{a'_i}^{a_i}(g_i(a_i, \cdot) - g_i(a'_i, \cdot)) - \gamma_{a'_i}^{a_i}(\cdot) - \delta_{a'_i}^{a_i}(\cdot) = -(g'_i(a_i, \cdot) - g'_i(a'_i, \cdot))$$

where

$$\gamma_{a'_i}^{a_i}(\cdot) = \sum_{a''_i \neq a_i, a'_i} u_{a''_i}^{a_i}(g_i(a_i, \cdot) - g_i(a''_i, \cdot)) + \sum_{a''_i \neq a_i, a'_i} v_{a''_i}^{a_i}(g_i(a'_i, \cdot) - g_i(a''_i, \cdot))$$

with $u_{a''_i}^{a_i}, v_{a''_i}^{a_i} \geq 0$ and

$$\delta_{a'_i}^{a_i}(\cdot) = \sum_{a_{-i} \in A_{-i}} z_{a_{-i}} \delta^{a_{-i}}(\cdot)$$

with $z_{a_{-i}} \geq 0$. Thus,

$$x_{a'_i}^{a_i}(g_i(a_i, \cdot) - g_i(a'_i, \cdot)) + \gamma_{a'_i}^{a_i}(\cdot) \leq g'_i(a_i, \cdot) - g'_i(a'_i, \cdot).$$

We show $x_{a'_i}^{a_i} > 0$. Suppose that $x_{a'_i}^{a_i} = 0$, i.e., $\gamma_{a'_i}^{a_i}(\cdot) \leq g'_i(a_i, \cdot) - g'_i(a'_i, \cdot)$. Let

$$\lambda'_i \in \Lambda_i(a_i, A_i \setminus \{a'_i\} | g_i) \setminus \Lambda_i(a_i, a'_i | g_i),$$

which exists by $a_i \approx_i^g a'_i$ and G3. Since $\lambda'_i \in \Lambda_i(a_i, A_i \setminus \{a'_i\} | g_i) \cap \Lambda_i(a'_i, A_i \setminus \{a_i\} | g_i)$,

$$\begin{aligned} & \sum_{a_{-i} \in A_{-i}} \lambda'_i(a_{-i}) \gamma_{a'_i}^{a_i}(a_{-i}) \\ &= \sum_{a''_i \neq a_i, a'_i} u_{a''_i}^{a_i} \sum_{a_{-i} \in A_{-i}} \lambda'_i(a_{-i}) (g_i(a_i, a_{-i}) - g_i(a''_i, a_{-i})) \\ &+ \sum_{a''_i \neq a_i, a'_i} v_{a''_i}^{a'_i} \sum_{a_{-i} \in A_{-i}} \lambda'_i(a_{-i}) (g_i(a'_i, a_{-i}) - g_i(a''_i, a_{-i})) \geq 0. \end{aligned}$$

Since $\lambda'_i \in \Lambda_i(a'_i, A_i | g_i) = \Lambda_i(a'_i, A_i | g'_i)$ and $\lambda'_i \notin \Lambda_i(a_i, A_i | g_i) = \Lambda_i(a_i, A_i | g'_i)$,

$$\sum_{a_{-i} \in A_{-i}} \lambda'_i(a_{-i}) (g'_i(a_i, a_{-i}) - g'_i(a'_i, a_{-i})) < 0.$$

This is a contradiction. Thus, we must have $x_{a'_i}^{a_i} > 0$.

We have

$$x_{a'_i}^{a_i} (g_i(a_i, \cdot) - g_i(a'_i, \cdot)) + \gamma_{a'_i}^{a_i}(\cdot) \leq g'_i(a_i, \cdot) - g'_i(a'_i, \cdot)$$

and symmetrically

$$x_{a_i}^{a'_i} (g_i(a'_i, \cdot) - g_i(a_i, \cdot)) + \gamma_{a_i}^{a'_i}(\cdot) \leq g'_i(a'_i, \cdot) - g'_i(a_i, \cdot)$$

where $x_{a'_i}^{a_i}, x_{a_i}^{a'_i} > 0$. Adding both,

$$(x_{a'_i}^{a_i} - x_{a_i}^{a'_i}) (g_i(a_i, \cdot) - g_i(a'_i, \cdot)) + \gamma_{a'_i}^{a_i}(\cdot) + \gamma_{a_i}^{a'_i}(\cdot) \leq 0. \tag{5}$$

We show $x_{a'_i}^{a_i} - x_{a_i}^{a'_i} = 0$. Suppose that $x_{a'_i}^{a_i} - x_{a_i}^{a'_i} > 0$. Let

$$\begin{aligned} \lambda_i & \in \Lambda_i(a'_i, A_i \setminus \{a_i\} | g_i) \setminus \Lambda_i(a'_i, a_i | g_i) \\ & \subseteq \Lambda_i(a_i, A_i \setminus \{a'_i\} | g_i) \cap \Lambda_i(a'_i, A_i \setminus \{a_i\} | g_i). \end{aligned}$$

Then, the expectation of the left-hand side of (5) is positive because

$$(x_{a'_i}^{a_i} - x_{a_i}^{a'_i}) \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i}) (g_i(a_i, a_{-i}) - g_i(a'_i, a_{-i})) > 0$$

and

$$\begin{aligned} & \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i}) (\gamma_{a'_i}^{a_i}(a_{-i}) + \gamma_{a_i}^{a'_i}(a_{-i})) \\ &= \sum_{a''_i \neq a_i, a'_i} (u_{a''_i}^{a_i} + v_{a''_i}^{a'_i}) \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i}) (g_i(a_i, a_{-i}) - g_i(a''_i, a_{-i})) \\ &+ \sum_{a''_i \neq a_i, a'_i} (v_{a''_i}^{a'_i} + u_{a''_i}^{a_i}) \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i}) (g_i(a'_i, a_{-i}) - g_i(a''_i, a_{-i})) \geq 0. \end{aligned}$$

This is a contradiction. Symmetrically, if $x_{a'_i}^{a_i} - x_{a_i}^{a'_i} < 0$, then we have the symmetric contradiction. Thus, $x_{a'_i}^{a_i} - x_{a_i}^{a'_i} = 0$, and (5) is reduced to

$$\gamma_{a'_i}^{a_i}(\cdot) + \gamma_{a_i}^{a'_i}(\cdot) \leq 0. \quad (6)$$

We show $\gamma_{a'_i}^{a_i}(\cdot) = \gamma_{a_i}^{a'_i}(\cdot) = 0$. Suppose that either $\gamma_{a'_i}^{a_i}(\cdot) \neq 0$ or $\gamma_{a_i}^{a'_i}(\cdot) \neq 0$ is true. Let $\lambda_i, \lambda'_i \in \Delta(A_{-i})$ be such that

$$\begin{aligned} \lambda_i &\in \Lambda_i(a'_i, A_i \setminus \{a_i\} \mid g_i) \setminus \Lambda_i(a'_i, a_i \mid g_i) \\ &\subseteq \Lambda_i(a_i, A_i \setminus \{a'_i\} \mid g_i) \cap \Lambda_i(a'_i, A_i \setminus \{a_i\} \mid g_i), \\ \lambda'_i &\in \Lambda_i(a_i, A_i \setminus \{a'_i\} \mid g_i) \setminus \Lambda_i(a_i, a'_i \mid g_i) \\ &\subseteq \Lambda_i(a_i, A_i \setminus \{a'_i\} \mid g_i) \cap \Lambda_i(a'_i, A_i \setminus \{a_i\} \mid g_i). \end{aligned}$$

Consider $(\lambda_i + \lambda'_i)/2 \in \Delta(A_{-i})$. Then, the expectation of the left-hand side of (6) is positive because

$$\begin{aligned} &\sum_{a_{-i} \in A_{-i}} \frac{\lambda_i(a_{-i}) + \lambda'_i(a_{-i})}{2} (\gamma_{a'_i}^{a_i}(a_{-i}) + \gamma_{a_i}^{a'_i}(a_{-i})) \\ &= \sum_{a''_i \neq a_i, a'_i} (u_{a''_i}^{a_i} + v_{a''_i}^{a'_i}) \sum_{a_{-i} \in A_{-i}} \frac{\lambda_i(a_{-i}) + \lambda'_i(a_{-i})}{2} (g_i(a_i, a_{-i}) - g_i(a'_i, a_{-i})) \\ &\quad + \sum_{a''_i \neq a_i, a'_i} (v_{a''_i}^{a'_i} + u_{a''_i}^{a_i}) \sum_{a_{-i} \in A_{-i}} \frac{\lambda_i(a_{-i}) + \lambda'_i(a_{-i})}{2} (g_i(a'_i, a_{-i}) - g_i(a_i, a_{-i})) \\ &\geq \sum_{a''_i \neq a_i, a'_i} (u_{a''_i}^{a_i} + v_{a''_i}^{a'_i}) \sum_{a_{-i} \in A_{-i}} \frac{\lambda_i(a_{-i})}{2} (g_i(a_i, a_{-i}) - g_i(a'_i, a_{-i})) \\ &\quad + \sum_{a''_i \neq a_i, a'_i} (v_{a''_i}^{a'_i} + u_{a''_i}^{a_i}) \sum_{a_{-i} \in A_{-i}} \frac{\lambda'_i(a_{-i})}{2} (g_i(a'_i, a_{-i}) - g_i(a_i, a_{-i})) > 0. \end{aligned}$$

This is a contradiction. Thus, $\gamma_{a'_i}^{a_i}(\cdot) = \gamma_{a_i}^{a'_i}(\cdot) = 0$.

Summarizing the above, we have

$$x_{a'_i}^{a_i}(g_i(a_i, \cdot) - g_i(a'_i, \cdot)) = g'_i(a_i, \cdot) - g'_i(a'_i, \cdot)$$

where $x_{a'_i}^{a_i} > 0$. This proves (b). \square

The following proposition and corollary follow by exactly the same arguments in Proposition 2 and Corollary 3 in the previous subsection for better-response equivalence.

Proposition 5. *Suppose that games \mathbf{g} and \mathbf{g}' satisfy generic properties G2 and G3, and that, for each $i \in N$ and for any $a_i, a'_i \in A_i$, there exists a sequence $\{a_i^k\}_{k=1}^m$ such that*

$a_i^1 = a_i, a_i^m = a'_i, a_i^k \approx_i^{\mathbf{g}} a_i^{k+1}$ for $k = 1, \dots, m - 1, a_i^k \approx_i^{\mathbf{g}} a_i^{k+2}$ for $k = 1, \dots, m - 2$. Then \mathbf{g} is best-response equivalent to \mathbf{g}' if and only if \mathbf{g} is VNM-equivalent to \mathbf{g}' .

Corollary 6. If \mathbf{g} and \mathbf{g}' satisfy generic properties G2 and G3 and $\approx_i^{\mathbf{g}}$ is the complete relation, then \mathbf{g} is best-response equivalent to \mathbf{g}' if and only if \mathbf{g} is VNM-equivalent to \mathbf{g}' .

4. Games with own-strategy unimodality

Best-response equivalence relation is an equivalence relation. It will be useful if, as a closed form, we can describe the best-response equivalence class of a game in which best-response equivalence is a strictly weaker requirement than VNM-equivalence.

Let A_i be linearly ordered such that $A_i = \{1, \dots, K_i\}$ with $K_i \geq 3$. For $q_i : A_{-i} \rightarrow \mathbb{R}$ and $w_i : A_i \setminus \{K_i\} \rightarrow \mathbb{R}_{++}$, let $(q_i, w_i) \circ g_i : A \rightarrow \mathbb{R}$ be such that

$$(q_i, w_i) \circ g_i(1, \cdot) = q_i(\cdot),$$

$$(q_i, w_i) \circ g_i(a_i, \cdot) = q_i(\cdot) + \sum_{k=1}^{a_i-1} w_i(k)(g_i(k+1, \cdot) - g_i(k, \cdot)) \quad \text{for } a_i \geq 2.$$

Let $\mathcal{D}_i(g_i)$ be a class of payoff functions of player i obtained by this transformation:

$$\mathcal{D}_i(g_i) = \{g'_i : A \rightarrow \mathbb{R} \mid g'_i = (q_i, w_i) \circ g_i, q_i : A_{-i} \rightarrow \mathbb{R}, w_i : A_i \setminus \{K_i\} \rightarrow \mathbb{R}_{++}\}.$$

It is straightforward to see that $g'_i \in \mathcal{D}_i(g_i)$ if and only if there exists $w_i : A_i \setminus \{K_i\} \rightarrow \mathbb{R}_{++}$ such that

$$g'_i(a_i + 1, \cdot) - g'_i(a_i, \cdot) = w_i(a_i)(g_i(a_i + 1, \cdot) - g_i(a_i, \cdot)) \tag{7}$$

for all $a_i \in A_i \setminus \{K_i\}$. Note that $g_i \in \mathcal{D}_i(g_i), g'_i \in \mathcal{D}_i(g_i)$ implies $g_i \in \mathcal{D}_i(g'_i)$, and $g'_i \in \mathcal{D}_i(g_i)$ with $g''_i \in \mathcal{D}_i(g'_i)$ implies $g''_i \in \mathcal{D}_i(g_i)$. Thus, $\mathcal{D}_i(g_i)$ defines an equivalence class of payoff functions of player i . We write

$$\mathcal{D}(\mathbf{g}) = \{\mathbf{g}' = (g'_i)_{i \in N} \mid g'_i \in \mathcal{D}_i(g_i) \text{ for all } i \in N\}.$$

For example, consider a parametrized class of games $\{\mathbf{g}(x, y)\}_{(x,y) \in \mathbb{R}_{++}^2}$ discussed in Section 2. We have that $\{\mathbf{g}(x, y)\}_{(x,y) \in \mathbb{R}_{++}^2} \subset \mathcal{D}(\mathbf{g}(1, 1))$. To see this, we write $\mathbf{g}(x, y) = (g_i(\cdot \mid x, y))_{i \in \{1,2\}}$. Then, for any $(x, y) \in \mathbb{R}_{++}^2$ and $i \neq j$,

$$g_i(1, a_j \mid x, y) = q_i(a_j),$$

$$g_i(2, a_j \mid x, y) = q_i(a_j) + x(g_i(2, a_j \mid 1, 1) - g_i(1, a_j \mid 1, 1)),$$

$$g_i(3, a_j \mid x, y) = q_i(a_j) + x(g_i(2, a_j \mid 1, 1) - g_i(1, a_j \mid 1, 1))$$

$$+ y(g_i(3, a_j \mid 1, 1) - g_i(2, a_j \mid 1, 1))$$

where $q_i : \{1, 2, 3\} \rightarrow \mathbb{R}$ is such that $q_i(1) = x, q_i(2) = -x$, and $q_i(3) = -2x$. Remember that, for any $(x, y) \in \mathbb{R}_{++}^2, \mathbf{g}(x, y)$ is best-response equivalent to $\mathbf{g}(1, 1)$. It is easy to see that every game in $\mathcal{D}(\mathbf{g}(1, 1))$ is VNM-equivalent to $\mathbf{g}(x, y)$ for some $(x, y) \in \mathbb{R}_{++}^2$. Thus, every game in $\mathcal{D}(\mathbf{g}(1, 1))$ is best-response equivalent to $\mathbf{g}(1, 1)$.

This observation leads us to the question when every game in $\mathcal{D}(\mathbf{g})$ is best-response equivalent to \mathbf{g} . We provide a necessary and sufficient condition for it.

We say that g_i is own-strategy unimodal if, for all $\lambda_i \in \Delta(A_{-i})$, there exists $k^* \in A_i$ such that,

$$\begin{aligned} \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(a_i, a_{-i}) - g_i(a_i - 1, a_{-i})) &\geq 0 \quad \text{if } a_i \leq k^* \text{ and } k^* > 1, \\ \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(a_i, a_{-i}) - g_i(a_i + 1, a_{-i})) &\geq 0 \quad \text{if } a_i \geq k^* \text{ and } k^* < K_i. \end{aligned} \tag{8}$$

Note that if g_i is own-strategy unimodal, then (8) is true if and only if $\lambda_i \in \Lambda_i(k^*, A_i \mid g_i)$. Clearly, by (7), g_i is own-strategy unimodal if and only if $g'_i \in \mathcal{D}_i(g_i)$ is own-strategy unimodal.

We say that g_i is own-strategy concave if $g_i(\cdot, a_{-i}) : A_i \rightarrow \mathbb{R}$ is concave, i.e., $g_i(a_i + 1, a_{-i}) - g_i(a_i, a_{-i})$ is decreasing in a_i for all $a_{-i} \in A_{-i}$.

Lemma 4. *Suppose that $g_i(a_i + 1, a_{-i}) \neq g_i(a_i, a_{-i})$ for all $a_i \in A_i \setminus \{K_i\}$ and $a_{-i} \in A_{-i}$, and that there is no weakly dominated strategy. Then, g_i is own-strategy unimodal if and only if there exists $\tilde{g}_i \in \mathcal{D}_i(g_i)$ such that \tilde{g}_i is own-strategy concave.*

Proof. Suppose that $\tilde{g}_i \in \mathcal{D}_i(g_i)$ is own-strategy concave. Then, $\tilde{g}_i(a_i + 1, a_{-i}) - \tilde{g}_i(a_i, a_{-i})$ is decreasing in a_i for all $a_{-i} \in A_{-i}$. Thus, $\sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(\tilde{g}_i(a_i + 1, a_{-i}) - \tilde{g}_i(a_i, a_{-i}))$ is also decreasing in a_i for all $\lambda_i \in \Delta(A_{-i})$. This immediately implies that $\tilde{g}_i \in \mathcal{D}_i(g_i)$ is own-strategy unimodal. Since

$$\begin{aligned} &\sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(a_i + 1, a_{-i}) - g_i(a_i, a_{-i})) \\ &= \frac{1}{w_i(a_i)} \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(\tilde{g}_i(a_i + 1, a_{-i}) - \tilde{g}_i(a_i, a_{-i})), \end{aligned}$$

g_i is also own-strategy unimodal.

Suppose that g_i is own-strategy unimodal. We prove the existence of an own-strategy concave payoff function $\tilde{g}_i = (q_i, w_i) \circ g_i$ by construction. Later, we will show that there exists $C_k > 0$ such that

$$g_i(k + 1, \cdot) - g_i(k, \cdot) \geq C_k(g_i(k + 2, \cdot) - g_i(k + 1, \cdot)). \tag{9}$$

For C_k satisfying (9), we let $w_i : A_i \rightarrow \mathbb{R}_{++}$ be such that $w_i(1) = 1$ and $w_i(a_i) = \prod_{k=1}^{a_i-1} C_k$ for $a_i \geq 2$, and $q_i : A_{-i} \rightarrow \mathbb{R}$ be such that $q_i(a_{-i}) = 0$ for all $a_{-i} \in A_{-i}$. Since

$$\tilde{g}_i(a_i + 1, \cdot) - \tilde{g}_i(a_i, \cdot) = w_i(a_i)(g_i(a_i + 1, \cdot) - g_i(a_i, \cdot)),$$

we have

$$\begin{aligned} \tilde{g}_i(k + 1, \cdot) - \tilde{g}_i(k, \cdot) &= w_i(k)(g_i(k + 1, \cdot) - g_i(k, \cdot)), \\ \tilde{g}_i(k + 2, \cdot) - \tilde{g}_i(k + 1, \cdot) &= C_k w_i(k)(g_i(k + 2, \cdot) - g_i(k + 1, \cdot)). \end{aligned}$$

By this and (9), we have

$$\tilde{g}_i(k + 1, \cdot) - \tilde{g}_i(k, \cdot) \geq \tilde{g}_i(k + 2, \cdot) - \tilde{g}_i(k + 1, \cdot),$$

which implies that \tilde{g}_i is own-strategy concave.

We prove the existence of C_k satisfying (9) by Farkas' Lemma. Before doing it, we must first observe that if

$$\sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(k + 1, a_{-i}) - g_i(k, a_{-i})) = 0 \tag{10}$$

then

$$\sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(k + 2, a_{-i}) - g_i(k + 1, a_{-i})) \leq 0.$$

To see this, suppose otherwise. Then, there exists $\lambda_i \in \Delta(A_{-i})$ satisfying both (10) and

$$\sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(k + 2, a_{-i}) - g_i(k + 1, a_{-i})) > 0.$$

Since $g_i(k + 1, a_{-i}) - g_i(k, a_{-i}) \neq 0$ for all $a_{-i} \in A_{-i}$, (10) implies that there exist $a'_{-i}, a''_{-i} \in A_{-i}$ such that $0 < \lambda_i(a'_{-i}) < 1$ with $g_i(k + 1, a'_{-i}) - g_i(k, a'_{-i}) > 0$ and $0 < \lambda_i(a''_{-i}) < 1$ with $g_i(k + 1, a''_{-i}) - g_i(k, a''_{-i}) < 0$. Let $\varepsilon > 0$ be sufficiently small. More precisely, let $\varepsilon > 0$ be such that

$$\varepsilon < \min \left\{ \lambda_i(a'_{-i}), 1 - \lambda_i(a''_{-i}), \frac{\sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(k + 2, a_{-i}) - g_i(k + 1, a_{-i}))}{2 \times \max_{a_{-i} \in A_{-i}} |g_i(k + 2, a_{-i}) - g_i(k + 1, a_{-i})|} \right\}.$$

Let $\lambda'_i \in \Delta(A_{-i})$ be such that

$$\lambda'_i(a_{-i}) = \begin{cases} \lambda_i(a_{-i}) - \varepsilon & \text{if } a_{-i} = a'_{-i}, \\ \lambda_i(a_{-i}) + \varepsilon & \text{if } a_{-i} = a''_{-i}, \\ \lambda_i(a_{-i}) & \text{otherwise.} \end{cases}$$

Then, we have

$$\begin{aligned} & \sum_{a_{-i} \in A_{-i}} \lambda'_i(a_{-i})(g_i(k + 1, a_{-i}) - g_i(k, a_{-i})) \\ &= \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(k + 1, a_{-i}) - g_i(k, a_{-i})) \\ & \quad + \varepsilon(g_i(k + 1, a''_{-i}) - g_i(k, a''_{-i})) - \varepsilon(g_i(k + 1, a'_{-i}) - g_i(k, a'_{-i})) \\ &= \varepsilon(g_i(k + 1, a''_{-i}) - g_i(k, a''_{-i})) - \varepsilon(g_i(k + 1, a'_{-i}) - g_i(k, a'_{-i})) < 0, \\ & \sum_{a_{-i} \in A_{-i}} \lambda'_i(a_{-i})(g_i(k + 2, a_{-i}) - g_i(k + 1, a_{-i})) \\ &= \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(k + 2, a_{-i}) - g_i(k + 1, a_{-i})) \\ & \quad + \varepsilon(g_i(k + 2, a''_{-i}) - g_i(k + 1, a''_{-i})) - \varepsilon(g_i(k + 2, a'_{-i}) - g_i(k + 1, a'_{-i})) \end{aligned}$$

$$\begin{aligned} &\geq \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(k+2, a_{-i}) - g_i(k+1, a_{-i})) \\ &\quad - 2\varepsilon \max_{a_{-i} \in A_{-i}} |g_i(k+2, a_{-i}) - g_i(k+1, a_{-i})| > 0, \end{aligned}$$

which contradicts to the assumption that g_i is own-strategy unimodal.

Now, we know that, if g_i is own-strategy unimodal and satisfies the assumptions, then it must be true that if

$$\sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(k+1, a_{-i}) - g_i(k, a_{-i})) \leq 0,$$

then

$$\sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(k+2, a_{-i}) - g_i(k+1, a_{-i})) \leq 0.$$

This implies that if $(y_{a_{-i}})_{a_{-i} \in A_{-i}} \in \mathbb{R}^{A_{-i}}$ is such that

$$\begin{aligned} &\sum_{a_{-i} \in A_{-i}} y_{a_{-i}}(g_i(k+1, a_{-i}) - g_i(k, a_{-i})) \leq 0, \\ &-y_{a_{-i}} \leq 0 \quad \text{for all } a_{-i} \in A_{-i}, \end{aligned}$$

then

$$\sum_{a_{-i} \in A_{-i}} y_{a_{-i}}(g_i(k+2, a_{-i}) - g_i(k+1, a_{-i})) \leq 0.$$

By Farkas' Lemma, there exist $x_k \geq 0$ and $z_{a_{-i}} \geq 0$ for $a_{-i} \in A_{-i}$ such that

$$x_k(g_i(k+1, \cdot) - g_i(k, \cdot)) - \sum_{a_{-i} \in A_{-i}} z_{a_{-i}} \delta^{a_{-i}}(\cdot) = g_i(k+2, \cdot) - g_i(k+1, \cdot).$$

Thus,

$$x_k(g_i(k+1, \cdot) - g_i(k, \cdot)) \geq g_i(k+2, \cdot) - g_i(k+1, \cdot). \quad (11)$$

If $x_k = 0$, then $g_i(k+2, \cdot) - g_i(k+1, \cdot) \leq 0$. However, this is impossible since there is no weakly dominated strategy. Thus, $x_k > 0$. By letting $C_k = 1/x_k$, (11) implies (9). \square

Consider again $\{\mathbf{g}(x, y)\}_{(x, y) \in \mathbb{R}_{++}^2} \subset \mathcal{D}(\mathbf{g}(1, 1))$. In general, $g_i(\cdot | x, y)$ is not always own-strategy concave. However, $g_i(\cdot | 1, 1)$ is own-strategy concave. Thus, Lemma 4 says that $g_i(\cdot | x, y)$ is own-strategy unimodal.

We claim that, generically, $\mathcal{D}(\mathbf{g})$ is a best-response equivalence class if and only if g_i is own-strategy unimodal for all $i \in N$.

Proposition 7. *Suppose that \mathbf{g} has no dominated strategy. Every game in $\mathcal{D}(\mathbf{g})$ is best-response equivalent to \mathbf{g} if and only if g_i is own-strategy unimodal for all $i \in N$. If g_i is own-strategy unimodal for all $i \in N$ and \mathbf{g} satisfies generic property G3, then every game best-response equivalent to \mathbf{g} and satisfying G3 is in $\mathcal{D}(\mathbf{g})$.*

Proof. Suppose that g_i is own-strategy unimodal for all $i \in N$. We show that if $\mathbf{g}' \in \mathcal{D}(\mathbf{g})$ then \mathbf{g}' is best-response equivalent to \mathbf{g} . Let $\lambda_i \in \Lambda_i(a_i^*, A_i | g_i)$. Then, (8) implies that

$$\begin{aligned} \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(a_i, a_{-i}) - g_i(a_i - 1, a_{-i})) &\geq 0 \quad \text{if } a_i \leq a_i^* \text{ and } a_i^* > 1, \\ \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(a_i, a_{-i}) - g_i(a_i + 1, a_{-i})) &\geq 0 \quad \text{if } a_i \geq a_i^* \text{ and } a_i^* < K_i. \end{aligned} \tag{12}$$

By (7), this is true if and only if

$$\begin{aligned} \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g'_i(a_i, a_{-i}) - g'_i(a_i - 1, a_{-i})) &\geq 0 \quad \text{if } a_i \leq a_i^* \text{ and } a_i^* > 1, \\ \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g'_i(a_i, a_{-i}) - g'_i(a_i + 1, a_{-i})) &\geq 0 \quad \text{if } a_i \geq a_i^* \text{ and } a_i^* < K_i. \end{aligned} \tag{13}$$

Thus, $\lambda_i \in \Lambda_i(a_i^*, A_i | g'_i)$. Conversely, let $\lambda_i \in \Lambda_i(a_i^*, A_i | g'_i)$. Since g'_i is own-strategy unimodal, we have (13), which is true if and only if (12) is true. Thus, $\lambda_i \in \Lambda_i(a_i^*, A_i | g_i)$. Therefore, $\Lambda_i(a_i^*, A_i | g_i) = \Lambda_i(a_i^*, A_i | g'_i)$ and thus \mathbf{g}' is best-response equivalent to \mathbf{g} .

Conversely, suppose that every game in $\mathcal{D}(\mathbf{g})$ is best-response equivalent to \mathbf{g} . We show that g_i is own-strategy unimodal for all $i \in N$. Seeking a contradiction, suppose otherwise. Then, there exist $a_i^*, \tilde{a}_i \in A_i$ and $\lambda_i \in \Lambda_i(a_i^*, A_i | g_i)$ such that either of the following is true:

$$a_i^* < \tilde{a}_i \quad \text{and} \quad \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(\tilde{a}_i, a_{-i}) - g_i(\tilde{a}_i - 1, a_{-i})) > 0, \tag{14}$$

$$a_i^* > \tilde{a}_i \quad \text{and} \quad \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(\tilde{a}_i, a_{-i}) - g_i(\tilde{a}_i + 1, a_{-i})) > 0. \tag{15}$$

When (14) is true, let $g'_i = (q_i, w_i) \circ g_i \in \mathcal{D}_i(g_i)$ be such that $q_i(\cdot) = 0$ and

$$w_i(a_i) = \begin{cases} L & \text{if } a_i = \tilde{a}_i - 1, \\ 1 & \text{otherwise.} \end{cases}$$

Then, we have

$$\begin{aligned} &\sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g'_i(\tilde{a}_i, a_{-i}) - g'_i(a_i^*, a_{-i})) \\ &= \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g'_i(\tilde{a}_i, a_{-i}) - g'_i(\tilde{a}_i - 1, a_{-i})) \\ &\quad + \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g'_i(\tilde{a}_i - 1, a_{-i}) - g'_i(a_i^*, a_{-i})) \\ &= L \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(\tilde{a}_i, a_{-i}) - g_i(\tilde{a}_i - 1, a_{-i})) \\ &\quad + \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(\tilde{a}_i - 1, a_{-i}) - g_i(a_i^*, a_{-i})). \end{aligned}$$

By choosing very large $L > 0$, we have

$$\sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g'_i(\tilde{a}_i, a_{-i}) - g'_i(a_i^*, a_{-i})) > 0$$

and thus $\Lambda_i(a_i^*, A_i | g_i) \neq \Lambda_i(a_i^*, A_i | g'_i)$. When (15) is true, we also have $\Lambda_i(a_i^*, A_i | g_i) \neq \Lambda_i(a_i^*, A_i | g'_i)$ by the similar argument. This implies that some game in $\mathcal{D}(\mathbf{g})$ is not best-response equivalent to \mathbf{g} , which completes the proof of the first half of the proposition.

We prove the last half of the proposition. Suppose that g_i is own-strategy unimodal for all $i \in N$ and that \mathbf{g} satisfies generic property G3. Let \mathbf{g}' be best-response equivalent to \mathbf{g} and satisfy G3. We show $\mathbf{g}' \in \mathcal{D}(\mathbf{g})$.

We first observe that $a_i \approx_i^{\mathbf{g}} a_i + 1$ for all $a_i \in A_i \setminus \{K_i\}$. To see this, let $\lambda_i^k \in \Lambda_i(k, A_i | g_i)$ for $k \in A_i$, which exists since \mathbf{g} has no dominated strategy. Note that if $\lambda_i = \lambda_i^k$ or $\lambda_i = \lambda_i^{k+1}$ then

$$\begin{aligned} \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})g_i(k, a_{-i}) &\geq \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})g_i(a_i, a_{-i}) && \text{for all } a_i \leq k, \\ \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})g_i(k+1, a_{-i}) &\geq \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})g_i(a_i, a_{-i}) && \text{for all } a_i \geq k+1. \end{aligned} \tag{16}$$

Let $t \in [0, 1]$ and $\lambda_i^{k,t} = t\lambda_i^k + (1-t)\lambda_i^{k+1} \in \Delta(A_{-i})$ be such that

$$\sum_{a_{-i} \in A_{-i}} \lambda_i^{k,t}(a_{-i})g_i(k, a_{-i}) = \sum_{a_{-i} \in A_{-i}} \lambda_i^{k,t}(a_{-i})g_i(k+1, a_{-i}). \tag{17}$$

Then, (16) implies that

$$\begin{aligned} \sum_{a_{-i} \in A_{-i}} \lambda_i^{k,t}(a_{-i})g_i(k, a_{-i}) &\geq \sum_{a_{-i} \in A_{-i}} \lambda_i^{k,t}(a_{-i})g_i(a_i, a_{-i}) && \text{for all } a_i \leq k, \\ \sum_{a_{-i} \in A_{-i}} \lambda_i^{k,t}(a_{-i})g_i(k+1, a_{-i}) &\geq \sum_{a_{-i} \in A_{-i}} \lambda_i^{k,t}(a_{-i})g_i(a_i, a_{-i}) && \text{for all } a_i \geq k+1. \end{aligned}$$

By (17), we have $\lambda_i^{k,t} \in \Lambda_i(k, A_i | g_i) \cap \Lambda_i(k+1, A_i | g_i)$. This implies that $a_i \approx_i^{\mathbf{g}} a_i + 1$ for all $a_i \in A_i \setminus \{K_i\}$.

Since \mathbf{g} and \mathbf{g}' satisfy G3 and are best-response equivalent, we can use Proposition 4, which says that there exists $w_i : A_i \setminus \{K_i\} \rightarrow \mathbb{R}_{++}$ such that

$$g'_i(a_i + 1, \cdot) - g'_i(a_i, \cdot) = w_i(a_i)(g_i(a_i + 1, \cdot) - g_i(a_i, \cdot)).$$

This implies that $g'_i \in \mathcal{D}_i(g_i)$ and thus $\mathbf{g}' \in \mathcal{D}(\mathbf{g})$. \square

A weaker, but similar claim is true for games such that strategy sets are intervals of real numbers and payoff functions are differentiable, which has a couple of applications. In the remainder of this section, we discuss this issue.

Abusing notation, we give a definition of best-response equivalence for a class of games with a continuum of actions. Let A_i be a closed interval of \mathbb{R} for all $i \in N$. Assume that $g_i : A \rightarrow \mathbb{R}$ is bounded and continuously differentiable. Let $\Delta(A_{-i})$ be the set of all probability measures over A_{-i} and $\Lambda_i(a_i, X_i | g_i)$ be such that

$$\Lambda_i(a_i, X_i \mid g_i) = \left\{ \lambda_i \in \Delta(A_{-i}) \mid \int_{A_{-i}} (g_i(a_i, a_{-i}) - g_i(a'_i, a_{-i})) d\lambda_i(a_{-i}) \geq 0 \text{ for all } a'_i \in X_i \right\}.$$

The definition of best-response equivalence is the same as that for finite games: we say that \mathbf{g} is best-response equivalent to \mathbf{g}' if, for each $i \in N$, $\Lambda_i(a_i, A_i \mid g_i) = \Lambda_i(a_i, A_i \mid g'_i)$ for all $a_i \in A_i$.

We say that g_i is own-strategy unimodal if, for any $\lambda_i \in \Delta(A_{-i})$, there exists x^* such that

$$\begin{aligned} \frac{\partial}{\partial a_i} \int_{A_{-i}} g_i(a_i, a_{-i}) d\lambda_i(a_{-i}) &\geq 0 \quad \text{if } a_i \leq x^* \text{ and } x^* > \min A_i, \\ \frac{\partial}{\partial a_i} \int_{A_{-i}} g_i(a_i, a_{-i}) d\lambda_i(a_{-i}) &\leq 0 \quad \text{if } a_i \geq x^* \text{ and } x^* < \max A_i. \end{aligned} \tag{18}$$

Note that if g_i is own-strategy unimodal, then (18) is true if and only if $\lambda_i \in \Lambda_i(x^*, A_i \mid g_i)$. Since

$$\frac{\partial}{\partial a_i} \int_{A_{-i}} g_i(a_i, a_{-i}) d\lambda_i(a_{-i}) = \int_{A_{-i}} \frac{\partial g_i(a_i, a_{-i})}{\partial a_i} d\lambda_i(a_{-i}),$$

g_i is own-strategy unimodal if g_i is own-strategy concave, i.e., $\partial g_i(a_i, a_{-i})/\partial a_i$ is decreasing in a_i for all $a_{-i} \in A_{-i}$.

For measurable functions $q_i : A_{-i} \rightarrow \mathbb{R}$ and $w_i : A_i \rightarrow \mathbb{R}_{++}$, let $(q_i, w_i) \circ g_i : A \rightarrow \mathbb{R}$ be such that, for $a_i \in A_i$ and $a_{-i} \in A_{-i}$,

$$(q_i, w_i) \circ g_i(a_i, a_{-i}) = q_i(a_{-i}) + \int_{x \leq a_i} w_i(x) \frac{\partial g_i(x, a_{-i})}{\partial x} dx.$$

Let

$$\begin{aligned} \mathcal{D}_i(g_i) &= \{g'_i : A \rightarrow \mathbb{R} \mid g'_i = (q_i, w_i) \circ g_i, q_i : A_{-i} \rightarrow \mathbb{R}, w_i : A_i \rightarrow \mathbb{R}_{++}\}, \\ \mathcal{D}(\mathbf{g}) &= \{\mathbf{g}' = (g'_i)_{i \in N} \mid g'_i \in \mathcal{D}_i(g_i)\}. \end{aligned}$$

Proposition 8. *Suppose that g_i is own-strategy unimodal for all $i \in N$. Then, every game in $\mathcal{D}(\mathbf{g})$ is best-response equivalent to \mathbf{g} .*

Proof. Let $\mathbf{g}' \in \mathcal{D}(\mathbf{g})$. Since g_i is own-strategy unimodal, for all $\lambda_i \in \Delta(A_i)$, there exists $a_i^* \in A_i$ such that

$$\begin{aligned} \frac{\partial}{\partial a_i} \int_{A_{-i}} g_i(a_i, a_{-i}) d\lambda_i(a_{-i}) &\geq 0 \quad \text{if } a_i \leq a_i^* \text{ and } a_i^* > \min A_i, \\ \frac{\partial}{\partial a_i} \int_{A_{-i}} g_i(a_i, a_{-i}) d\lambda_i(a_{-i}) &\leq 0 \quad \text{if } a_i \geq a_i^* \text{ and } a_i^* < \max A_i. \end{aligned} \tag{19}$$

Since

$$\frac{\partial g'_i(a_i, a_{-i})}{\partial a_i} = w_i(a_i) \frac{\partial g_i(a_i, a_{-i})}{\partial a_i},$$

(19) is true if and only if

$$\begin{aligned} \frac{\partial}{\partial a_i} \int_{A_{-i}} g'_i(a_i, a_{-i}) d\lambda_i(a_{-i}) &\geq 0 \quad \text{if } a_i \leq a_i^* \text{ and } a_i^* > \min A_i, \\ \frac{\partial}{\partial a_i} \int_{A_{-i}} g'_i(a_i, a_{-i}) d\lambda_i(a_{-i}) &\leq 0 \quad \text{if } a_i \geq a_i^* \text{ and } a_i^* < \max A_i. \end{aligned} \quad (20)$$

Thus, g'_i is also own-strategy unimodal. Since (19) is true if and only if $\lambda_i \in \Lambda_i(a_i^*, A_i | g_i)$ and (20) is true if and only if $\lambda_i \in \Lambda_i(a_i^*, A_i | g'_i)$, we must have $\Lambda_i(a_i^*, A_i | g_i) = \Lambda_i(a_i^*, A_i | g'_i)$, which completes the proof. \square

This proposition has a useful application concerning the uniqueness of correlated equilibria. Neyman (1997) showed that if \mathbf{g} has a continuously differentiable and strictly concave potential function,⁹ then the potential maximizer is the unique correlated equilibrium of \mathbf{g} . The set of correlated equilibria is the same for two games if the two games are best-response equivalent. Thus, we claim the following.

Corollary 9. *Suppose that \mathbf{g} has a continuously differentiable and strictly concave potential function f . Then, the potential maximizer is the unique correlated equilibrium of every game in $\mathcal{D}(\mathbf{g})$.*

Note that a game in $\mathcal{D}(\mathbf{g})$ is not necessarily a potential game and payoff functions are not necessarily concave.

5. Mixed extensions of equivalence

We have focused on players' preferences over pure strategies, given nondegenerate conjectures about their opponents' behavior. But we could ask the same question in the mixed strategy extension of the original game; equivalently, we could look at players' preferences over mixed strategies.¹⁰ The natural question is whether or not our discussion so far must be modified by the "mixed extension" of equivalence.

For $i \in N$, let $\Delta(A_i)$ denote the set of all mixed strategies of player i . Abusing notation, we write $g_i(p_i, a_{-i}) = \sum_{a_i \in A_i} p_i(a_i) g_i(a_i, a_{-i})$ for $p_i \in \Delta(A_i)$. By the mixed extension of Λ_i , we can naturally define $\Lambda_i(p_i, X_i | g_i)$ for $p_i \in \Delta(A_i)$ and $X_i \subseteq \Delta(A_i)$:

$$\Lambda_i(p_i, X_i | g_i)$$

⁹ The definition of potential functions of this class of games is the same as those of finite games.

¹⁰ The associate editor suggested the observations in this section.

$$= \left\{ \lambda_i \in \Delta(A_{-i}) \mid \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(p_i, a_{-i}) - g_i(p'_i, a_{-i})) \geq 0 \text{ for all } p'_i \in X_i \right\}.$$

We use the same rule $\Lambda_i(p_i, p'_i \mid g_i) = \Lambda_i(p_i, \{p'_i\} \mid g_i)$ as before. We consider the following equivalence relations of games.

Definition 7. A game \mathbf{g} is *mixed better-response equivalent* to \mathbf{g}' if, for each $i \in N$,

$$\Lambda_i(p_i, p'_i \mid g_i) = \Lambda_i(p_i, p'_i \mid g'_i)$$

for all $p_i, p'_i \in \Delta(A_i)$.

Definition 8. A game \mathbf{g} is *mixed best-response equivalent* to \mathbf{g}' if, for each $i \in N$,

$$\Lambda_i(p_i, \Delta(A_i) \mid g_i) = \Lambda_i(p_i, \Delta(A_i) \mid g'_i)$$

for all $p_i \in \Delta(A_i)$.

Note that VNM-equivalence is sufficient for both mixed better-response equivalence and mixed best-response equivalence. Note also that mixed better-response equivalence is sufficient for better-response equivalence, and that mixed best-response equivalence is sufficient for best-response equivalence. It is easy to see that mixed best-response equivalence is not only sufficient but also necessary for best-response equivalence.

Lemma 5. A game \mathbf{g} is *mixed best-response equivalent* to \mathbf{g}' if and only if \mathbf{g} is *best-response equivalent* to \mathbf{g}' .

Proof. Note that

$$\begin{aligned} & \lambda_i \in \Lambda_i(p_i, \Delta(A_i) \mid g_i) \\ \Leftrightarrow & p_i \in \arg \max_{p'_i \in \Delta(A_i)} \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})g_i(p'_i, a_{-i}) \\ \Leftrightarrow & p_i(a_i) > 0 \text{ implies } a_i \in \arg \max_{a'_i \in A_i} \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})g_i(a'_i, a_{-i}) \\ \Leftrightarrow & p_i(a_i) > 0 \text{ implies } \lambda_i \in \Lambda_i(a_i, A_i \mid g_i). \end{aligned}$$

Thus, if $\Lambda_i(a_i, A_i \mid g_i) = \Lambda_i(a_i, A_i \mid g'_i)$ for all $a_i \in A_i$, then $\Lambda_i(p_i, \Delta(A_i) \mid g_i) = \Lambda_i(p_i, \Delta(A_i) \mid g'_i)$ for all $p_i \in \Delta(A_i)$. This completes the proof. \square

This lemma implies that the characterization of mixed best-response equivalence is reduced to that of best-response equivalence.

On the other hand, mixed better-response equivalence is a strictly stronger requirement than better-response equivalence. Consider a two player, three strategy, symmetric payoff games \mathbf{g} and \mathbf{g}' , where each player's payoffs are given by the following payoff matrices

(where the player’s own strategies are represented by rows and his opponent’s strategies are represented by columns):

	g		
	1	2	3
1	1	-2	-2
2	0	0	0
3	2	-1	-1

	g'		
	1	2	3
1	x	-2x	-2x
2	0	0	0
3	2y	-y	-y

We assume that $x, y > 0$ and $x/2 < y < 2x$. Then, $1 \sim_i^g 2, 2 \sim_i^g 3, 3 \succ_i^g 1$, and $\succ_i^g = \succ_i^{g'}$. We also have $x(g_i(1, \cdot) - g_i(2, \cdot)) = g'_i(1, \cdot) - g'_i(2, \cdot)$ and $y(g_i(2, \cdot) - g_i(3, \cdot)) = g'_i(2, \cdot) - g'_i(3, \cdot)$. Thus, by Proposition 1, **g** is better-response equivalent to **g'**. However, we can show that **g** is mixed better-response equivalent to **g'** only if $x = y$. To see this, suppose that a row player believes that the column player never chooses 1: a row player has a belief λ_i with $\lambda_i(1) = 0$. Consider a row player’s mixed strategy p_i such that $p_i(1) = p$ and $p_i(2) = 1 - p$. In **g**, he prefers strategy p_i to strategy 3 if and only if $-2p \geq -1$, i.e., $p \leq 1/2$. In **g'**, he prefers strategy p_i to strategy 3 if and only if $-2xp \geq -y$, i.e., $p \leq y/2x$. In order for **g** to be mixed better-response equivalent to **g'**, it must be true that $1/2 = y/2x$, i.e., $x = y$. In this case, **g** is VNM-equivalent to **g'** and thus mixed better-response equivalent to **g'**.

In the above example, the relation \sim_i^g generates a connected graph since $1 \sim_i^g 2$ and $2 \sim_i^g 3$. Thus, under the connectedness of \sim_i^g , better-response equivalence does not necessarily imply VNM-equivalence, but mixed better-response equivalence may imply VNM-equivalence. The natural question is whether this is true. Remember that Proposition 2 provides a condition to ensure the equivalence of better-response equivalence and VNM-equivalence. The condition includes the connectedness of \sim_i^g . But the connectedness is not sufficient as demonstrated by the above example. In contrast, the following proposition asserts that the connectedness of \sim_i^g ensures the equivalence of mixed better-response equivalence and VNM-equivalence.

Proposition 10. *Suppose that games **g** and **g'** satisfy generic properties G1 and G2, and that, for each $i \in N$, \sim_i^g generates a connected graph on A_i . Then **g** is mixed better-response equivalent to **g'** if and only if **g** is VNM-equivalent to **g'**.*

To prove the proposition, we use the following lemma.

Lemma 6. *Suppose that **g** and **g'** satisfy generic properties G1 and G2, and that **g** is mixed better-response equivalent to **g'**. For distinct $a_i, b_i, c_i \in A_i$, if $a_i \sim_i^g b_i$ and $b_i \sim_i^g c_i$, then there exists $w_i > 0$ such that*

$$g_i(a_i, \cdot) - g_i(b_i, \cdot) = w_i(g'_i(a_i, \cdot) - g'_i(b_i, \cdot)),$$

$$g_i(b_i, \cdot) - g_i(c_i, \cdot) = w_i(g'_i(b_i, \cdot) - g'_i(c_i, \cdot)).$$

Proof. By G1 and Proposition 1, there exist $w_i(a_i, b_i), w_i(b_i, c_i) > 0$ such that

$$g_i(a_i, \cdot) - g_i(b_i, \cdot) = w_i(a_i, b_i)(g'_i(a_i, \cdot) - g'_i(b_i, \cdot)),$$

$$g_i(b_i, \cdot) - g_i(c_i, \cdot) = w_i(b_i, c_i)(g'_i(b_i, \cdot) - g'_i(c_i, \cdot)).$$

We show that $w_i(a_i, b_i) = w_i(b_i, c_i)$.

Either $a_i \sim_i^g c_i$, $a_i \succ_i^g c_i$, or $c_i \succ_i^g a_i$ is true. If $a_i \sim_i^g c_i$, there exists $w_i(a_i, c_i) > 0$ such that

$$g_i(a_i, \cdot) - g_i(c_i, \cdot) = w_i(a_i, c_i)(g'_i(a_i, \cdot) - g'_i(c_i, \cdot))$$

by Proposition 1. Thus, by Lemma 3, $w_i(a_i, b_i) = w_i(b_i, c_i) = w_i(a_i, c_i)$.

Suppose that $a_i \succ_i^g c_i$. Note that $\succ_i^g = \succ_i^{g'}$ by Proposition 1. Let $\lambda_i \in \Delta(A_{-i})$ be such that

$$\sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g'_i(b_i, a_{-i}) - g'_i(c_i, a_{-i})) < 0, \tag{21}$$

which exists since $b_i \sim_i^{g'} c_i$ and G1. The relation $a_i \succ_i^{g'} c_i$ implies that

$$\sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g'_i(a_i, a_{-i}) - g'_i(c_i, a_{-i})) > 0. \tag{22}$$

By the weighted average of (21) and (22), we can choose $p_i \in \Delta(A_i)$ such that $p_i(a_i) = p$, $p_i(b_i) = 1 - p$, and

$$\sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g'_i(p_i, a_{-i}) - g'_i(c_i, a_{-i})) = 0. \tag{23}$$

Mixed better-response equivalence implies that

$$\sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g_i(p_i, a_{-i}) - g_i(c_i, a_{-i})) = 0. \tag{24}$$

Now calculate

$$\begin{aligned} g_i(p_i, \cdot) - g_i(c_i, \cdot) &= pg_i(a_i, \cdot) + (1 - p)g_i(b_i, \cdot) - g_i(c_i, \cdot) \\ &= p(g_i(a_i, \cdot) - g_i(b_i, \cdot)) + g_i(b_i, \cdot) - g_i(c_i, \cdot) \\ &= w_i(a_i, b_i)p(g'_i(a_i, \cdot) - g'_i(b_i, \cdot)) \\ &\quad + w_i(b_i, c_i)(g'_i(b_i, \cdot) - g'_i(c_i, \cdot)) \\ &= w_i(a_i, b_i)(pg'_i(a_i, \cdot) + (1 - p)g'_i(b_i, \cdot) - g'_i(c_i, \cdot)) \\ &\quad + (w_i(b_i, c_i) - w_i(a_i, b_i))(g'_i(b_i, \cdot) - g'_i(c_i, \cdot)) \\ &= w_i(a_i, b_i)(g'_i(p_i, \cdot) - g'_i(c_i, \cdot)) \\ &\quad + (w_i(b_i, c_i) - w_i(a_i, b_i))(g'_i(b_i, \cdot) - g'_i(c_i, \cdot)). \end{aligned}$$

By the expectations with respect to λ_i for both sides of the equation, and by (23) and (24), we have

$$(w_i(b_i, c_i) - w_i(a_i, b_i)) \sum_{a_{-i} \in A_{-i}} \lambda_i(a_{-i})(g'_i(b_i, a_{-i}) - g'_i(c_i, a_{-i})) = 0.$$

By (21), we must have $w_i(a_i, b_i) = w_i(b_i, c_i)$. Similarly, if $c_i \succ_i^g a_i$, we must have $w_i(a_i, b_i) = w_i(b_i, c_i)$. This completes the proof. \square

We now report the proof of Proposition 10.

Proof of Proposition 10. We show that if \mathbf{g} is mixed better-response equivalent to \mathbf{g}' then \mathbf{g} is VNM-equivalent to \mathbf{g}' . By G1 and Proposition 1, if $a_i \sim_i^g a'_i$, there exists $w_i(a_i, a'_i) > 0$ such that

$$g_i(a_i, \cdot) - g_i(a'_i, \cdot) = w_i(a_i, a'_i)(g'_i(a_i, \cdot) - g'_i(a'_i, \cdot)).$$

If $|A_i| = 2$, this completes the proof by Lemma 1. Suppose that $|A_i| \geq 3$. For $a_i, a'_i \in A_i$, let $\{a_i^k\}_{k=1}^m$ be a sequence such that $a_i^1 = a_i$, $a_i^m = a'_i$, $a_i^k \sim_i^g a_i^{k+1}$ for $k = 1, \dots, m-1$, which exists by the connectedness of \sim_i^g . There exists $x_k > 0$ such that

$$g_i(a_i^k, \cdot) - g_i(a_i^{k+1}, \cdot) = x_k(g'_i(a_i^k, \cdot) - g'_i(a_i^{k+1}, \cdot)).$$

By Lemma 6, $x_k = x_{k+1}$ for all $k \leq m-1$. By letting $x_k = w_i(a_i, a'_i)$, we have

$$\begin{aligned} g_i(a_i, \cdot) - g_i(a'_i, \cdot) &= \sum_{k=1}^{m-1} (g_i(a_i^k, \cdot) - g_i(a_i^{k+1}, \cdot)) \\ &= \sum_{k=1}^{m-1} x_k (g'_i(a_i^k, \cdot) - g'_i(a_i^{k+1}, \cdot)) \\ &= w_i(a_i, a'_i) (g'_i(a_i, \cdot) - g'_i(a'_i, \cdot)). \end{aligned}$$

To summarize, for all $a_i, a'_i \in A_i$, there exists $w_i(a_i, a'_i) > 0$ satisfying the above equation. By Lemma 3, $w_i(a_i, a'_i)$ is the same for all $a_i, a'_i \in A_i$. By Lemma 1, \mathbf{g} is VNM-equivalent to \mathbf{g}' , which completes the proof. \square

Acknowledgments

We are very grateful for valuable input from Larry Blume, George Mailath and Philip Reny. Ui acknowledges financial support by the Ministry of Education, Science, Sports and Culture, Grant-in-Aid for Scientific Research.

References

- Anderson, S.P., Jacob, K., Holt, C.A., 2001. Minimum-effort coordination games: stochastic potential and logit equilibrium. *Games Econ. Behav.* 34, 177–199.
- Blume, L., 1993. The statistical mechanics of strategic interaction. *Games Econ. Behav.* 5, 387–424.
- Brock, W., Durlauf, S., 2001. Discrete choice with social interactions. *Rev. Econ. Stud.* 68, 235–260.
- Dubey, P., Haimanko, O., Zapechelnyuk, A., 2002. Strategic substitutes and potential games. Mimeo. SUNY at Stony Brook.
- Hiriart-Urruty, J.-B., Lemaréchal, C., 2001. *Fundamentals of Convex Analysis*. Springer-Verlag, New York.
- Maskin, E., Tirole, J., 2001. Markov perfect equilibrium. *J. Econ. Theory* 100, 191–219.

- Mertens, J.-F., 1987. Ordinality in non-cooperative games. Mimeo. DP8728, CORE, Université Catholique de Louvain.
- Monderer, D., Shapley, L.S., 1996a. Fictitious play property for games with identical interests. *J. Econ. Theory* 68, 258–265.
- Monderer, D., Shapley, L.S., 1996b. Potential games. *Games Econ. Behav.* 14, 124–143.
- Morris, S., 1999. Potential methods in interaction games. Mimeo. Yale University. Available from <http://www.econ.yale.edu/~sm326/pot-method.pdf>.
- Morris, S., Ui, T., 2002. Generalized potentials and robust sets of equilibria. Mimeo. Yale University. Available from <http://www.econ.yale.edu/~sm326/genpot.pdf>.
- Neyman, A., 1997. Correlated equilibrium and potential games. *Int. J. Game Theory* 26, 223–227.
- Rockafellar, R.T., 1970. *Convex Analysis*. Princeton Univ. Press, Princeton, NJ.
- Rosenthal, R.W., 1973. A class of games possessing pure strategy equilibria. *Int. J. Game Theory* 2, 65–67.
- Sela, A., 1992. Learning processes in games. MSc thesis. The Technion, Haifa, Israel. [In Hebrew.]
- Sela, A., 1999. Fictitious play in ‘one-against-all’ multi-player games. *Econ. Theory* 14, 635–651.
- Ui, T., 2000. A Shapley value representation of potential games. *Games Econ. Behav.* 31, 121–135.
- Ui, T., 2001. Robust equilibria of potential games. *Econometrica* 69, 1373–1380.
- Ui, T., 2002. Quantal response equilibria and stochastic best response dynamics. Mimeo. Yokohama National University. Available from <http://www2.igss.ynu.ac.jp/~oui/qresse.pdf>.
- Voorneveld, M., 2000. Best-response potential games. *Econ. Letters* 66, 289–295.